



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



PII: S0049-3848(20)30130-4

DOI: <https://doi.org/10.1016/j.thromres.2020.04.014>

Reference: TR 7653

To appear in: *Thrombosis Research*

Received date: 11 April 2020

Accepted date: 13 April 2020

Please cite this article as: R. Escher, N. Breakey and B. Lämmle, Severe COVID-19 infection associated with endothelial activation, *Thrombosis Research* (2018), <https://doi.org/10.1016/j.thromres.2020.04.014>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Severe COVID-19 infection associated with endothelial activation

Robert Escher¹, Neal Breakey¹, Bernhard Lämmle^{2,3,4}¹ Department of Medicine, Spital Emmental, Burgdorf, Switzerland² Department of Hematology and Central Hematology Laboratory, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland³ Center for Thrombosis and Hemostasis, University Medical Center Mainz, Germany⁴ Haemostasis Research Unit, University College London, London, UK

Mounting evidence supports the strong prognostic importance of D-dimers and the impact of coagulopathy in COVID-19 patients^{1,2,3}. We would like to emphasize the significance of anticoagulation in severe COVID-19 disease, by adding our observation of highly pathological data on anti-phospholipid-antibodies, von Willebrand Factor (VWF) and Factor VIII. We observed a 72-year-old previously healthy male patient admitted to our hospital 6 days after the onset of respiratory symptoms and fever. Six days after admission, following a rapid deterioration of his clinical condition with development of acute respiratory distress syndrome, acute renal insufficiency and altered mental status, he was transferred to the intensive care unit (ICU). Supportive care with intubation, positive pressure ventilation and renal replacement therapy led to a stabilization of his condition. During the ICU stay, a continual increase of D-dimers was observed, from initially 0.69 mg/L to 2.55 mg/L on day 4, 2.81 mg/L on day 11 to 20.63 mg/L on day 21. At this time, IgG anti-cardiolipin antibodies (ACA) and anti-beta2-glycoprotein I (anti- β 2-GPI) were negative, but IgM ACA elevated at 121.9 CU (normal <20 CU) and IgM anti- β 2-GPI elevated at 275.3 CU (normal <20 CU). Furthermore, on day 21, we also observed a concurrent massive elevation of VWF, with VWF:antigen 555% (normal 42-136%) and VWF:activity 520% (normal 42-168%), accompanied by an increase of Factor VIII clotting activity of 369% (normal 55-164%). The increased VWF points towards massive endothelial stimulation and damage with release of VWF from Weibel-

Palade bodies. Interestingly, endothelial cells express ACE2, the receptor for SARS-CoV-2, thus possibly mediating endothelial activation³. The patient received prophylactic anticoagulation with dalteparin 5000 U s.c. daily from admission to hospital. We increased the dose to therapeutic anticoagulation with unfractionated heparin on day 21, without hemorrhagic complication. Since then, the patient is steadily improving, ventilation and hemodialysis could be successfully withdrawn, and he has been transferred from the ICU to a general ward. D-dimers decreased to 6.26 mg/L on day 24 and to 1.94 mg/L on day 29. Considering the high levels of VWF and factor VIII, we strongly suggest administering higher, possibly therapeutic doses of anticoagulation to these patients. The role of antiphospholipid antibodies, VWF and FVIII needs further study.

References

1. F.A. Klok, M.J.H.A. Kruip, N.J.M. van der Meer, M.S. Arbous, D.A.M.P.J. Gommers, K.M. Kant, F.H.J. Kaptein, J. van Paassen, M.A.M. Stals, M.V. Huisman, H. Endeman, Incidence of thrombotic complications in critically ill ICU patients with COVID-19, *Thromb. Res.* (n.d.). <https://doi.org/10.1016/j.thromres.2020.04.013>.
2. F. Zhou, T. Yu, R. Du, G. Fan, Y. Liu, Z. Liu, J. Xiang, Y. Wang, B. Song, X. Gu, L. Guan, Y. Wei, H. Li, X. Wu, J. Xu, S. Tu, Y. Zhang, H. Chen, B. Cao, Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study, *Lancet.* 0 (2020). [https://doi.org/10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3).
3. Y. Zhang, M. Xiao, S. Zhang, P. Xia, W. Cao, W. Jiang, H. Chen, X. Ding, H. Zhao, H. Zhang, C. Wang, J. Zhao, X. Sun, R. Tian, W. Wu, D. Wu, J. Ma, Y. Chen, D. Zhang, J. Xie, X. Yan, X. Zhou, Z. Liu, J. Wang, B. Du, Y. Qin, P. Gao, X. Qin, Y. Xu, W. Zhang, T. Li, F. Zhang, Y. Zhao, Y. Li, S. Zhang, Coagulopathy and Antiphospholipid Antibodies in Patients with Covid-19, *N. Engl. J. Med.* (2020). <https://doi.org/10.1056/NEJMc2007575>.
4. I. Hamming, W. Timens, M.L.C. Bulthuis, A.T. Lely, G.J. Navis, H. van Goor, Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis, *J. Pathol.* 203 (2004) 631–637.